

After Bedside *Removal* of a Peritoneal Dialysis Catheter: What do I need to know?

1. Dressing care

- DO NOT change your dressing yourself. Your PD nurse will change/remove your dressing at your follow-up appointment.
- Keep your dressing clean and dry at all times.
- If your dressing becomes loose, add more tape.
- If your dressing becomes wet or loose, put an extra dressing on top. If the extra dressing becomes wet or bloody, call your kidney nurse or doctor.
- 2. No showers, bathing or swimming. Sponge baths only until your kidney nurse or doctor tells you otherwise (usually after your stitches are taken out).

3. If you have pain

- You may take acetaminophen (Tylenol®).
- A little bit of pain is normal but it should decrease each day. If the pain gets worse, talk with your kidney nurse or doctor.



4. Attend your follow-up appointment

- Usually one to two weeks after your catheter has been removed.
- Your stitches and the dressing will be removed.

Date and time of follow-up appointment

5. When should I call my kidney nurse or doctor?

- Your extra dressing (the one that was put on top of the original dressing) gets wet or bloody.
- Your pain will not go away even after pain medications
- Your temperature is more than 38.0° C (100.4° F)

Important phone numbers

Kidney Doctor _____

Peritoneal Dialysis Unit ____

Kidney Clinic _

The information in this pamphlet is provided for educational and information purposes, and to support discussion with your health care team about your medical condition and treatment. It does not constitute medical advice and should not substitute advice given by your physician or other qualified health care professional.

